



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ :C07D 221/12, A61K 31/435, C07D
491/04 // (C07D 491/04, 317:00, 221:00)

A1

(11) International Publication Number:

WO 99/05112

(43) International Publication Date:

4 February 1999 (04.02.99)

(21) International Application Number: PCT/EP98/04476

(22) International Filing Date: 18 July 1998 (18.07.98)

(30) Priority Data:

97112794.9

25 July 1997 (25.07.97)

EP

(71) Applicant (for all designated States except US): BYK
GULDEN LOMBERG CHEMISCHE FABRIK GMBH
[DE/DE]; Byk-Gulden-Strasse 2, D-78467 Konstanz (DE).(72) Inventors (for all designated States except CA US): AM-
SCHLER, Hermann; Hohenhewenstrasse 19, D-78315
Radolfzell (DE). FLOCKERZI, Dieter; Ackerweg 26,
D-78476 Allensbach (DE). ULRICH, Wolf-Rüdiger;
Hebelstrasse 3, D-78464 Konstanz (DE). BÄR, Thomas;
Blarerstrasse 16, D-78462 Konstanz (DE). MARTIN,
Thomas; Sonnenbühlstrasse 73, D-78464 Konstanz
(DE). SCHUDT, Christian; Schützenstrasse 20, D-78462
Konstanz (DE). HATZELMANN, Armin; Alter Wall 3,
D-78467 Konstanz (DE). BEUME, Rolf; Bohlstrasse 13,
D-78465 Konstanz (DE). HÄFNER, Dietrich; Beethoven-
strasse 5, D-78464 Konstanz (DE). BOSS, Hildegard;
Flurweg 3a, D-78464 Konstanz (DE). KLEY, Hans-Peter;
Im Weinberg 3b, D-78476 Allensbach (DE).


(72) Inventor; and

(75) Inventor/Applicant (for US only): GUTTERER, Beate
[DE/DE]; Allensbacher Strasse 6b, D-78476 Allensbach
(DE).(74) Common Representative: BYK GULDEN LOMBERG
CHEMISCHE FABRIK GMBH; Byk-Gulden-Strasse 2,
D-78467 Konstanz (DE).(81) Designated States: AL, AU, BA, BG, BR, CA, CN, CZ, EE,
GE, HR, HU, ID, IL, JP, KR, LT, LV, MK, MX, NO, NZ,
PL, RO, SG, SI, SK, TR, UA, US, VN, YU, ZW, Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR,
IE, IT, LU, MC, NL, PT, SE).

Published

With international search report.

(54) Title: SUBSTITUTED 6-ALKYLPHENANTHRIDINES



(57) Abstract

Compounds of formula (I) in which R1, R2, R3, R31, R4, R5, R51, A and R6 have the meanings indicated in the description, are novel efficacious bronchial therapeutics.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

SUBSTITUTED 6-ALKYLPHENANTHRIDINES

Field of application of the invention

The invention relates to novel 6-alkylphenanthridines which are used in the pharmaceutical industry for the production of medicaments.


Known technical background

German Patent Application DE 11 99 773 describes the preparation of 6-methyl-8,9-methylenedioxy-1,2,3,4,4a,5,6,10b-octahydrophenanthridine, which should have analgesic, antiinflammatory and antipyretic properties. In the Indian Journal of Chemistry 7, 1969, 674-677 the preparation of ethyl 1,2,3,4,4a,10b-hexahydrophenanthridine-6-butyrate is described.

Description of the invention

It has now been found that the novel 6-alkylphenanthridines which are described below in greater detail have surprising and particularly advantageous properties.

The invention thus relates to compounds of the formula I



in which

R1 is hydroxyl, 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or completely or predominantly fluorine-substituted 1-4C-alkoxy,

R2 is hydroxyl, 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or completely or predominantly fluorine-substituted 1-4C-alkoxy,

or in which

R1 and R2 together are a 1-2C-alkylenedioxy group,

R3 is hydrogen or 1-4C-alkyl,

R31 is hydrogen or 1-4C-alkyl,

or in which

R3 and R31 together are a 1-4C-alkylene group,

R4 is hydrogen or 1-4C-alkyl,

R5 is hydrogen,

R51 is hydrogen,

or in which

R5 and R51 together are an additional bond,

A is 1-6C-alkylene or 5-7C-cycloalkylene,

R6 is COOR₆₁ or CON(R₆₂)R₆₃, where

R₆₁ is hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl or 3-7C-cycloalkylmethyl and

R₆₂ and R₆₃ independently of one another are hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl or 3-7C-cycloalkylmethyl,

and the salts of these compounds.

1-4C-Alkyl represents a straight-chain or branched alkyl radical having 1 to 4 carbon atoms. Examples which may be mentioned are the butyl, isobutyl, sec-butyl, tert-butyl, propyl, isopropyl and, preferably, the ethyl and methyl radicals.

1-4C-Alkoxy represents radicals which, in addition to the oxygen atom, contain a straight-chain or branched alkyl radical having 1 to 4 carbon atoms. Examples which may be mentioned are the butoxy, isobutoxy, sec-butoxy, tert-butoxy, propoxy, isopropoxy and, preferably, the ethoxy and methoxy radicals.

3-7C-Cycloalkoxy represents cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy and cycloheptyloxy, of which cyclopropyloxy, cyclobutyloxy and cyclopentyloxy are preferred.

3-7C-Cycloalkylmethoxy represents cyclopropylmethoxy, cyclobutylmethoxy, cyclopentylmethoxy, cyclohexylmethoxy and cycloheptylmethoxy, of which cyclopropylmethoxy, cyclobutylmethoxy and cyclopentylmethoxy are preferred.

Completely or predominantly fluorine-substituted 1-4C-alkoxy which may be mentioned are, for example, the 2,2,3,3,3-pentafluoropropoxy, the perfluoroethoxy, the 1,2,2-trifluoroethoxy, in particular the 1,1,2,2-tetrafluoroethoxy, the trifluoromethoxy, the 2,2,2-trifluoroethoxy and, preferably, the difluoro-

romethoxy radicals. "Predominantly" in this connection means that more than half of the hydrogen atoms are substituted by fluorine atoms.

1-2C-Alkylenedioxy represents, for example, the methylenedioxy ($-O-CH_2-O-$) and the ethylenedioxy radicals ($-O-CH_2-CH_2-O-$).

If R3 and R31 together have the meaning 1-4C-alkylene, the positions 1 and 4 in compounds of the formula I are linked to one another by a 1-4C-alkylene bridge, 1-4C-alkylene representing straight-chain or branched alkylene radicals having 1 to 4 carbon atoms. Examples which may be mentioned are the radicals methylene ($-CH_2-$), ethylene ($-CH_2-CH_2-$), trimethylene ($-CH_2-CH_2-CH_2-$), 1,2-dimethylethylene [$-CH(CH_3)-CH(CH_3)-$] and isopropylidene [$-C(CH_3)_2-$].

If R5 and R51 together are an additional bond, then the carbon atoms in the positions 2 and 3 in compounds of the formula I are linked to one another via a double bond.

1-6C-Alkylene represents straight-chain or branched 1-6C-alkylene radicals, for example the methylene ($-CH_2-$), ethylene ($-CH_2-CH_2-$), trimethylene ($-CH_2-CH_2-CH_2-$), tetramethylene ($-CH_2-CH_2-CH_2-CH_2-$), 1,2-dimethylethylene [$-CH(CH_3)-CH(CH_3)-$], 1,1-dimethylethylene [$-C(CH_3)_2-CH_2-$], 2,2-dimethylethylene [$-CH_2-C(CH_3)_2-$], isopropylidene [$-C(CH_3)_2-$], 1-methylethylene [$-CH(CH_3)-CH_2-$], pentamethylene ($-CH_2-CH_2-CH_2-CH_2-CH_2-$) and the hexamethylene radicals ($-CH_2-CH_2-CH_2-CH_2-CH_2-CH_2-$).

5-7C-Cycloalkylene represents cycloalkylene radicals having 5 to 7 carbon atoms. Cyclohexylene radicals are preferred, where, for example, the 1,3- and the 1,4-cyclohexylene radicals may be mentioned.

1-7C-Alkyl represents straight-chain or branched alkyl radicals having 1 to 7 carbon atoms. Examples which may be mentioned are the heptyl, isoheptyl (5-methylhexyl), hexyl, isohexyl (4-methylpentyl), neoheptyl (3,3-dimethylbutyl), pentyl, isopentyl (3-methylbutyl), neopentyl (2,2-dimethylpropyl), butyl, isobutyl, sec-butyl, tert-butyl, propyl, isopropyl, ethyl and methyl radicals.

3-7C-Cycloalkyl represents the cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl radicals.

3-7C-Cycloalkylmethyl represents a methyl radical which is substituted by one of the abovementioned 3-7C-cycloalkyl radicals. Examples which may be mentioned are the cyclopentylmethyl and the cyclohexylmethyl radicals.

Suitable salts for compounds of the formula I - depending on substitution - are all acid addition salts or all salts with bases. Particular mention may be made of the pharmacologically tolerable salts of the inorganic and organic acids and bases customarily used in pharmacy. Those which are suitable are, on the one hand, water-soluble and water-insoluble acid addition salts with acids such as, for example, hydrochloric acid, hydrobromic acid, phosphoric acid, nitric acid, sulfuric acid, acetic acid, citric acid, D-gluconic acid, benzoic acid, 2-(4-hydroxybenzoyl)benzoic acid, butyric acid, sulfosalicylic acid, maleic acid, lauric acid, malic acid, fumaric acid, succinic acid, oxalic acid, tartaric acid, embonic acid, stearic acid, toluenesulfonic acid, methanesulfonic acid or 3-hydroxy-2-naphthoic acid, where the acids are employed in salt preparation - depending on whether it is a mono- or polybasic acid and depending on which salt is desired, in an equimolar quantitative ratio or one differing therefrom.

On the other hand, salts with bases are also suitable. Examples of salts with bases which may be mentioned are alkali metal (lithium, sodium, potassium) or calcium, aluminum, magnesium, titanium, ammonium, meglumine or guanidinium salts, where here too the bases are employed in salt preparation in an equimolar quantitative ratio or one differing therefrom.

Pharmacologically intolerable salts which may be obtained initially as process products, for example in the preparation of the compounds according to the invention on an industrial scale, are converted into pharmacologically tolerable salts by processes known to the person skilled in the art.

According to expert's knowledge the compounds of the invention as well as their salts may contain, e.g. when isolated in crystalline form, varying amounts of solvents. Included within the scope of the invention are therefore all solvates and in particular all hydrates of the compounds of formula I as well as all solvates and in particular all hydrates of the salts of the compounds of formula I.

Compounds of the formula I to be emphasized are those in which

R1 is 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or completely or predominantly fluorine-substituted 1-2C-alkoxy,

R2 is 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or completely or predominantly fluorine-substituted 1-2C-alkoxy,

R3 is hydrogen,

R31 is hydrogen,

or in which

R3 and R31 together are a 1-2C-alkylene group,

R4 is hydrogen or 1-4C-alkyl,

R5 is hydrogen,

R51 is hydrogen,

or in which

R5 and R51 together are an additional bond,

A is 1-6C-alkylene or 5-7C-cycloalkylene,
R6 is COOR61 or CON(R62)R63, where
R61 is hydrogen, 1-7C-alkyl or 3-7C-cycloalkylmethyl and
R62 and R63 independently of one another are hydrogen or 1-7C-alkyl,
and the salts of these compounds.

Compounds of the formula I particularly to be emphasized are those in which

R1 is 1-4C-alkoxy, 3-7C-cycloalkoxy or completely or predominantly fluorine-substituted 1-2C-alkoxy,
R2 is 1-4C-alkoxy, 3-7C-cycloalkoxy or completely or predominantly fluorine-substituted 1-2C-alkoxy,
R3 is hydrogen,
R31 is hydrogen,

or in which

R3 and R31 together are a 1-2C-alkylene group,

R4 is hydrogen or 1-4C-alkyl,
R5 is hydrogen,
R51 is hydrogen,

or in which

R5 and R51 together are an additional bond,

A is 1-6C-alkylene or 5-7C-cycloalkylene,
R6 is COOR61 or CON(R62)R63, where
R61 is hydrogen, 1-4C-alkyl and
R62 and R63 independently of one another are hydrogen or 1-4C-alkyl,
and the salts of these compounds.


Preferred compounds of the formula I are those in which

R1 is 1-2C-alkoxy,
R2 is 1-2C-alkoxy,
R3 is hydrogen,
R31 is hydrogen,
R4 is hydrogen,
R5 is hydrogen,
R51 is hydrogen,
A is cyclohexylene,
R6 is COOR61, where
R61 is 1-4C-alkyl,

and the salts of these compounds.


The compounds of the formula I are chiral compounds having chiral centers in the positions 4a and 10b and, depending on the meaning of the substituents R3, R31, R4, R5 and R51, further chiral centers in the positions 1, 2, 3 and 4. If A represents 5-7C-cycloalkylene, depending on the regiochemistry of the substitution, 2 further chiral centers also occur within the alkylene ring. Also, if A is 1-6C-alkylene and the 1-6C-alkylene radical is branched, additional chiral centers can occur within the alkylene radical.

Numbering:



The invention therefore includes all conceivable pure diastereomers and pure enantiomers and their mixtures in any mixing ratio, including the racemates. The compounds of the formula I are preferred in which the hydrogen atoms in the positions 4a and 10b are cis to one another. Particularly preferred here are the pure cis diastereomers and the pure cis enantiomers and their mixtures in any mixing ratio and including the racemates.

The enantiomers can be separated in a manner known per se (for example by preparation and separation of appropriate diastereoisomeric compounds). Preferably, a separation of enantiomers takes place at the stage of the starting compounds of the formula III




for example via salt formation of the racemic compounds of the formula III with optically active carboxylic acids. Alternatively, enantiomerically pure starting compounds of the formula III can also be prepared via asymmetric syntheses.

If in the compounds of the formula I A is 5-7C-cycloalkylene or a branched 1-6C-alkylene radical, the stereochemistry within the alkylene radical is determined by the stereochemistry of the starting alkylene compounds used for the synthesis.

The invention further relates to a process for the preparation of the compounds of the formula I, in which R1, R2, R3, R31, R4, R5, R51, A and R6 have the meanings indicated above, and their salts.

The process comprises cyclocondensing compounds of the formula II



in which R1, R2, R3, R31, R4, R5, R51, A and R6 have the meanings indicated above, and, if desired, then converting the compounds of the formula I obtained into their salts, or, if desired, then converting salts of the compounds of the formula I obtained into the free compounds.


If desired, compounds of the formula I obtained can be converted into further compounds of the formula I by derivatization. For example, the corresponding acids can be obtained from compounds of the formula I, in which R6 is an ester group, by acidic or alkaline hydrolysis, the corresponding amides can be prepared by reaction with amines of the formula $\text{HN}(\text{R}_{62})\text{R}_{63}$ or alternatively corresponding esters can be prepared by transesterification of esters of the formula I or by esterification of acids of the formula I. The reactions are expediently carried out analogously to the methods known to the person skilled in the art.

Cyclocondensation is carried out in a manner known per se to the person skilled in the art according to Bischler-Napieralski (e.g. as described in J. Chem. Soc., 1956, 4280-4282) in the presence of a suitable condensing agent, such as, for example, polyphosphoric acid, phosphorus pentachloride, phosphorus trichloride, phosphorus pentoxide, thionyl chloride or preferably phosphorus oxychloride, in a suitable inert solvent, e.g. in a chlorinated hydrocarbon such as chloroform, or in a cyclic hydrocarbon such as toluene or xylene, or another inert solvent such as acetonitrile, or without further solvent using an excess of condensing agent, preferably at elevated temperature, in particular at the boiling temperature of the solvent or condensing agent used.

Compounds of the formula II, in which R1, R2, R3, R31, R4, R5, R51, A and R6 have the meanings indicated above, are accessible from the corresponding compounds of the formula III, in which R1, R2, R3, R31, R4, R5 and R51 have the meanings indicated above, by reaction with compounds of the formula R6-A-CO-X, in which R6 and A have the meanings indicated above and X is a suitable leaving group, preferably a chlorine atom. For example, benzylation is carried out as in the following examples according to the Einhorn process, the Schotten-Baumann variant or as described in J. Chem. Soc. (C), 1971, 1805-1808.

Compounds of the formula R6-A-CO-X and compounds of the formula III are either known or can be prepared in a known manner.

The compounds of the formula III can be prepared, for example, from compounds of the formula IV



in which R1, R2, R3, R31, R4, R5 and R51 have the meanings mentioned above, by reduction of the nitro group.


Reduction is carried out in a manner known to the person skilled in the art, for example as described in J. Org. Chem. 1962, 27, 4426 or as described in the following examples. Preferably, reduction is carried out by catalytic hydrogenation, e.g. in the presence of Raney nickel, in a lower alcohol such as methanol or ethanol at room temperature and under normal or elevated pressure. If desired, a catalytic amount of an acid, such as, for example, hydrochloric acid, can be added to the solvent.

The compounds of the formula III, in which R1, R2, R3, R31 and R4 have the meanings indicated above and R5 and R51 together are an additional bond, can be prepared from the corresponding compounds of the formula IV by selective reduction of the nitro group in a manner known to the person skilled in the art, for example in the presence of Raney nickel in a lower alcohol as a solvent using hydrazine hydrate as a hydrogen donor.

The compounds of the formula IV, in which R1, R2, R3, R31 and R4 have the meanings indicated above and R5 and R51 are hydrogen, are either known or can be prepared from corresponding compounds of the formula IV, in which R5 and R51 together are an additional bond. The reaction can be

carried out in a manner known to the person skilled in the art, preferably by hydrogenation in the presence of a catalyst, such as, for example, palladium on activated carbon, e.g. as described in J. Chem. Soc. (C), 1971, 1805-1808.

The compounds of the formula IV, in which R5 and R51 together are an additional bond, are either known or can be obtained by reaction of compounds of the formula V



in which R1 and R2 have the abovementioned meanings, with compounds of the formula VI




in which R3, R31 and R4 have the abovementioned meanings.

Cycloaddition is carried out here in a manner known to the person skilled in the art according to Diels-Alder, e.g. as described in J. Amer. Chem. Soc. 1957, 79, 6559 or in J. Org. Chem. 1952, 17, 581 or as described in the following examples.

Compounds of the formula IV obtained in the cycloaddition, in which the phenyl ring and the nitro group are trans to one another, can be converted into the corresponding cis compounds in a manner known to the person skilled in the art, e.g. as described in J. Amer. Chem. Soc. 1957, 79, 6559 or as described in the following examples.

The compounds of the formulae V and VI are either known or can be prepared in a known manner. The compounds of the formula V can be prepared, for example, from corresponding compounds of the formula VII in a manner known to the person skilled in the art, as described, for example, in J. Chem. Soc. 1951, 2524 or in J. Org. Chem. 1944, 9, 170 or as described in the following examples.

The compounds of the formula VII



in which R1 and R2 have the meanings indicated above, are either known or can be prepared in a manner known to the person skilled in the art, as described, for example, in Ber. Dtsch. Chem. Ges. **1925**, 58, 203.

The isolation and purification of the substances according to the invention is carried out in a manner known per se, for example by distilling off the solvent in vacuo and recrystallizing the resulting residue from a suitable solvent or subjecting it to one of the customary purification methods, such as, for example column chromatography on suitable support material.

Salts are obtained by dissolving the free compound in a suitable solvent, e.g. in a chlorinated hydrocarbon, such as methylene chloride or chloroform, or a low molecular weight aliphatic alcohol (ethanol, isopropanol), which contains the desired acid or base, or to which the desired acid or base is then added. The salts are obtained by filtering, reprecipitation, precipitation with a nonsolvent for the addition salt or by evaporation of the solvent. Salts obtained can be converted by alkalization or by acidification into the free compounds, which in turn can be converted into salts. In this manner, pharmacologically intolerable salts can be converted into pharmacologically tolerable salts.

The following examples serve to explain the invention in greater detail without restricting it. Likewise, further compounds of the formula I whose preparation is not described explicitly can be prepared in an analogous manner or in a manner familiar to the person skilled in the art using customary process techniques.

In the examples, m.p. stands for melting point, h for hour(s), RT for room temperature, EF for empirical formula, MW for molecular weight, calc. for calculated, fnd for found. The compounds and their salts mentioned in the examples are a preferred subject of the invention.

Examples

Final products

1. (+/-)-cis-(4a,10b)-9-Ethoxy-8-methoxy-6-(trans-4-ethoxycarbonylcyclohexyl)-1,2,3,4,4a,10b-hexahydrophenanthridine

3.0 g of methyl (+/-)-trans-4-[cis-(1,2)-2-(3-ethoxy-4-methoxyphenyl)cyclohexyl-1-carbamoyl]cyclohexane-1-carboxylate (compound A1) are dissolved in 100 ml of acetonitrile and 3.0 ml of phosphorus oxychloride and stirred overnight at 80°C. The reaction mixture is treated with 60 ml of ethyl acetate and extracted with sodium hydrogencarbonate solution. The organic phase is dried using sodium sulfate and concentrated. 2.0 g of the title compound are obtained as a solidifying oil.

EF: C₂₄H₃₃NO₄; MW 399.53

Elemental analysis:	calc.:	C 72.15 H 8.33 N 3.51
	find:	C 72.07 H 8.22 N 3.42

Starting compounds

- A1. Methyl (+/-)-trans-4-[cis-(1,2)-2-(3-ethoxy-4-methoxyphenyl)cyclohexyl-1-carbamoyl]cyclohexane-1-carboxylate

3.0 g of (+/-)-cis-2-ethoxy-1-methoxy-4-(2-aminocyclohexyl)benzene (compound B1) are dissolved in 50 ml of methylene chloride and 10 ml of triethylamine. A solution of 2.95 g of methyl trans-4-chlorocarbonylcyclohexanecarboxylate is added dropwise at RT and the mixture is extracted after stirring overnight with 100 ml each of water, 2 N hydrochloric acid, saturated sodium hydrogencarbonate solution and water again. The organic phase is dried using sodium sulfate and concentrated. The residue is extracted by stirring with ethyl acetate/petroleum ether in the ratio 7/3, filtered off with suction and dried. 3.8 g of the title compound of m.p. 147-148°C are obtained.

- B1. (+/-)-cis-2-Ethoxy-1-methoxy-4-(2-aminocyclohexyl)benzene

40.0 g of (+/-)-cis-2-ethoxy-1-methoxy-4-(2-nitrocyclohex-4-enyl)benzene (compound C1) are dissolved in 1000 ml of ethanol and 500 ml of tetrahydrofuran, treated with 10 g of Raney nickel and hydrogenated at a hydrogen pressure of 100 bar for 4 days in an autoclave. After filtration and removal of the solvent in vacuo, 35.9 g of the title compound are obtained as a solidifying oil.

C1. (+/-)-cis-2-Ethoxy-1-methoxy-4-(2-nitrocyclohex-4-enyl)benzene

89.25 g of (+/-)-trans-2-ethoxy-1-methoxy-4-(2-nitrocyclohex-4-enyl)benzene (compound D1) and 37 g of potassium hydroxide are dissolved in 500 ml of absolute ethanol. A solution of 23.5 ml of conc. sulfuric acid in 120 ml of absolute ethanol is then added dropwise such that the internal temperature does not exceed -2°C. After stirring for 1 h, the mixture is added to 4 l of ice water, and the precipitate is filtered off with suction, washed with water and dried. M.p.: 66-67°C.

D1. (+/-)-trans-2-Ethoxy-1-methoxy-4-(2-nitrocyclohex-4-enyl)benzene

110 g of 3-ethoxy-2-methoxy- ω -nitrostyrene (compound E1) and 360 mg of hydroquinone are suspended in 360 ml of absolute toluene and treated at -70°C with 180 ml of liquid 1,3-butadiene. The mixture is stirred at 160-180°C for 6 days in an autoclave and then cooled. The product is extracted by stirring with ethanol, filtered off with suction and dried. M.p.: 130-131°C.

E1. 3-Ethoxy-2-methoxy- ω -nitrostyrene

236 g of 3-ethoxy-2-methoxybenzaldehyde, 101 g of ammonium acetate and 320 ml of nitromethane are heated to 100°C for 4 h in 655 ml of glacial acetic acid. The solution is added to 5 l of ice water, and the precipitate is filtered off with suction, washed with water and dried. M.p.: 132-133°C.

Commercial utility

The compounds according to the invention have useful pharmacological properties which make them industrially utilizable. As selective cyclic nucleotide phosphodiesterase (PDE) inhibitors (specifically of type 4), they are suitable on the one hand as bronchial therapeutics (for the treatment of airway obstructions on account of their dilating action but also on account of their respiratory rate- or respiratory drive-increasing action) and for the removal of erectile dysfunction on account of their vascular dilating action, but on the other hand especially for the treatment of disorders, in particular of an inflammatory nature, e.g. of the airways (asthma prophylaxis), of the skin, of the intestine, of the eyes, of the CNS and of the joints, which are mediated by mediators such as histamine, PAF (platelet-activating factor), arachidonic acid derivatives such as leukotrienes and prostaglandins, cytokines, interleukins, chemokines, alpha-, beta- and gamma-interferon, tumor necrosis factor (TNF) or oxygen free radicals and proteases. In this context, the compounds according to the invention are distinguished by a low toxicity, a good enteral absorption (high bioavailability), a large therapeutic breadth and the absence of significant side effects.

On account of their PDE-inhibiting properties, the compounds according to the invention can be employed in human and veterinary medicine as therapeutics, where they can be used, for example, for the treatment and prophylaxis of the following illnesses: acute and chronic (in particular inflammatory and allergen-induced) airway disorders of varying origin (bronchitis, allergic bronchitis, bronchial asthma); dermatoses (especially of proliferative, inflammatory and allergic type) such as psoriasis (vulgaris), toxic and allergic contact eczema, atopic eczema, seborrhoeic eczema, Lichen simplex, sunburn, pruritus in the anogenital area, alopecia areata, hypertrophic scars, discoid lupus erythematosus, follicular and widespread pyodermias, endogenous and exogenous acne, acne rosacea and other proliferative, inflammatory and allergic skin disorders; disorders which are based on an excessive release of TNF and leukotrienes, for example disorders of the arthritis type (rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis and other arthritic conditions), disorders of the immune system (AIDS, multiple sclerosis), types of shock (septic shock, endotoxin shock, gram-negative sepsis, toxic shock syndrome and ARDS (adult respiratory distress syndrome)) and also generalized inflammations in the gastrointestinal region (Crohn's disease and ulcerative colitis); disorders which are based on allergic and/or chronic, immunological false reactions in the region of the upper airways (pharynx, nose) and the adjacent regions (paranasal sinuses, eyes), such as allergic rhinitis/sinusitis, chronic rhinitis/sinusitis, allergic conjunctivitis and also nasal polyps; but also disorders of the heart which can be treated by PDE inhibitors, such as cardiac insufficiency, or disorders which can be treated on account of the tissue-relaxant action of the PDE inhibitors, such as, for example, erectile dysfunction or colics of the kidneys and of the ureters in connection with kidney stones; and also illnesses of the central nervous system, such as depressions or arteriosclerotic dementia.

The invention further relates to a method for the treatment of mammals, including humans, which are suffering from one of the abovementioned illnesses. The method is characterized in that a therapeutically active and pharmacologically effective and tolerable amount of one or more of the compounds according to the invention is administered to the ill mammal.

The invention further relates to the compounds according to the invention for use in the treatment and/or prophylaxis of illnesses, especially the illnesses mentioned.

The invention also relates to the use of the compounds according to the invention for the production of medicaments which are employed for the treatment and/or prophylaxis of the illnesses mentioned.

The invention furthermore relates to medicaments for the treatment and/or prophylaxis of the illnesses mentioned, which contain one or more of the compounds according to the invention.

The medicaments are prepared by processes which are known per se and familiar to the person skilled in the art. As medicaments, the compounds according to the invention (= active compounds) are either employed as such, or preferably in combination with suitable pharmaceutical auxiliaries, e.g. in the form of tablets, coated tablets, capsules, suppositories, patches, emulsions, suspensions, gels or solutions, the active compound content advantageously being between 0.1 and 95%.

The person skilled in the art is familiar with auxiliaries which are suitable for the desired pharmaceutical formulations on account of his expert knowledge. In addition to solvents, gel formers, ointment bases and other active compound excipients, for example antioxidants, dispersants, emulsifiers, preservatives, solubilizers or permeation promoters, can be used.

For the treatment of disorders of the respiratory tract, the compounds according to the invention are preferably also administered by inhalation. To do this, these are either administered directly as a powder (preferably in micronized form) or by atomizing solutions or suspensions which contain them. With respect to the preparations and administration forms, reference is made, for example, to the details in European Patent 163 965.

For the treatment of dermatoses, the compounds according to the invention are in particular administered in the form of those medicaments which are suitable for topical application. For the production of the medicaments, the compounds according to the invention (= active compounds) are preferably mixed with suitable pharmaceutical auxiliaries and further processed to give suitable pharmaceutical formulations. Suitable pharmaceutical formulations are, for example, powders, emulsions, suspensions, sprays, oils, ointments, fatty ointments, creams, pastes, gels or solutions.

The medicaments according to the invention are prepared by processes known per se. The dosage of the active compounds is carried out in the order of magnitude customary for PDE inhibitors. Topical application forms (such as ointments) for the treatment of dermatoses thus contain the active com-

pounds in a concentration of, for example, 0.1-99%. The dose for administration by inhalation is customarily between 0.1 and 3 mg per day. The customary dose in the case of systemic therapy (p.o. or i.v.) is between 0.03 and 3 mg/kg per day.

Biological investigations

In the investigation of PDE 4 inhibition on the cellular plane, the activation of inflammatory cells is ascribed particular importance. An example is FMLP (N-formyl-methionyl-leucyl-phenylalanine)-induced superoxide production of neutrophilic granulocytes, which can be measured as luminol-amplified chemiluminescence. (Mc Phail LC, Strum SL, Leone PA and Sozzani S, The neutrophil respiratory burst mechanism. In "Immunology Series" 57: 47-76, 1992; ed. Coffey RG (Marcel Decker, Inc., New York-Basel-Hong Kong)).

Substances which inhibit chemiluminescence and cytokine secretion and the secretion of proinflammatory mediators on inflammatory cells, in particular neutrophilic and eosinophilic granulocytes, T-lymphocytes, monocytes and macrophages are those which inhibit PDE 4. This isoenzyme of the phosphodiesterase families is particularly represented in granulocytes. Its inhibition leads to an increase in the intracellular cyclic AMP concentration and thus to the inhibition of cellular activation. PDE 4 inhibition by the substances according to the invention is thus a central indicator for the suppression of inflammatory processes. (Giembycz MA, Could isoenzyme-selective phosphodiesterase inhibitors render bronchodilatory therapy redundant in the treatment of bronchial asthma?. *Biochem Pharmacol* 43: 2041-2051, 1992; Torphy TJ et al., Phosphodiesterase inhibitors: new opportunities for treatment of asthma. *Thorax* 46: 512-523, 1991; Schudt C et al., Zardaverine: a cyclic AMP PDE 3/4 inhibitor. In "New Drugs for Asthma Therapy", 379-402, Birkhäuser Verlag Basel 1991; Schudt C et al., Influence of selective phosphodiesterase inhibitors on human neutrophil functions and levels of cAMP and Ca; *Naunyn-Schmiedeberg's Arch Pharmacol* 344: 682-690, 1991; Tenor H and Schudt C, Analysis of PDE isoenzyme profiles in cells and tissues by pharmacological methods. In "Phosphodiesterase Inhibitors", 21-40, "The Handbook of Immunopharmacology", Academic Press, 1996; Hatzelmann A et al., Enzymatic and functional aspects of dual-selective PDE3/4-Inhibitors. In "Phosphodiesterase Inhibitors", 147-160, "The Handbook of Immunopharmacology", Academic Press, 1996.

1. Inhibition of PDE 4 activity**Methodology**

The activity test was carried out by the method of Bauer and Schwabe, which was adapted to micro-titer plates (Naunyn-Schmiedeberg's Arch. Pharmacol. 1980, 311, 193-198). In this connection, the PDE reaction is carried out in the first step. In a second step, the resultant 5'-nucleotide is cleaved to give the uncharged nucleoside by a 5'-nucleotidase of the snake venom from *Crotalus atrox*. In the third step, the nucleoside is separated from the remaining charged substrate on ion exchange columns. The columns are eluted with 2 ml of 30 mM ammonium formate (pH 6.0) directly into minivials to which 2 ml of scintillation fluid is additionally added for counting.


The inhibitory values determined for the compounds according to the invention [inhibitory concentration as $-\log IC_{50}$ (mol/l)] follow from Table A below, in which the numbers of the compounds correspond to the numbers of the examples.

Table A**Inhibition of PDE 4 activity**

Compound	$-\log IC_{50}$
1	7.47

Patent claims

1. A compound of the formula I



in which

R1 is hydroxyl, 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or completely or predominantly fluorine-substituted 1-4C-alkoxy,

R2 is hydroxyl, 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or completely or predominantly fluorine-substituted 1-4C-alkoxy,

or in which

R1 and R2 together are a 1-2C-alkylenedioxy group,

R3 is hydrogen or 1-4C-alkyl,

R31 is hydrogen or 1-4C-alkyl,

or in which

R3 and R31 together are a 1-4C-alkylene group,

R4 is hydrogen or 1-4C-alkyl,

R5 is hydrogen,

R51 is hydrogen,

or in which

R5 and R51 together are an additional bond,

A is 1-6C-alkylene or 5-7C-cycloalkylene,

R6 is COOR61 or CON(R62)R63, where

R61 is hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl or 3-7C-cycloalkylmethyl and

R62 and R63 independently of one another are hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl or 3-7C-cycloalkylmethyl,

or the salts of this compound.

2. A compound of the formula I as claimed in claim 1, in which

R1 is 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or completely or predominantly fluorine-substituted 1-2C-alkoxy,

R2 is 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or completely or predominantly fluorine-substituted 1-2C-alkoxy,

R3 is hydrogen,

R31 is hydrogen,

or in which

R3 and R31 together are a 1-2C-alkylene group,

R4 is hydrogen or 1-4C-alkyl,

R5 is hydrogen,

R51 is hydrogen,

or in which

R5 and R51 together are an additional bond,

A is 1-6C-alkylene or 5-7C-cycloalkylene,

R6 is COOR₆₁ or CON(R₆₂)R₆₃, where

R₆₁ is hydrogen, 1-7C-alkyl or 3-7C-cycloalkylmethyl and

R₆₂ and R₆₃ independently of one another are hydrogen or 1-7C-alkyl,

or the salts of this compound.

3. A compound of the formula I as claimed in claim 1, in which

R1 is 1-4C-alkoxy, 3-7C-cycloalkoxy or completely or predominantly fluorine-substituted 1-2C-alkoxy,

R2 is 1-4C-alkoxy, 3-7C-cycloalkoxy or completely or predominantly fluorine-substituted 1-2C-alkoxy,

R3 is hydrogen,

R31 is hydrogen,

or in which

R3 and R31 together are a 1-2C-alkylene group,

R4 is hydrogen or 1-4C-alkyl,

R5 is hydrogen,

R51 is hydrogen,

or in which

R5 and R51 together are an additional bond,

A is 1-6C-alkylene or 5-7C-cycloalkylene,

R6 is COOR₆₁ or CON(R₆₂)R₆₃, where

R₆₁ is hydrogen, 1-4C-alkyl and

R₆₂ and R₆₃ independently of one another are hydrogen or 1-4C-alkyl,

or the salts of this compound.

4. A compound of the formula I as claimed in claim 1, in which

R1 is 1-2C-alkoxy,

R2 is 1-2C-alkoxy,

R3 is hydrogen,

R31 is hydrogen,

R4 is hydrogen,

R5 is hydrogen,

R51 is hydrogen,

A is cyclohexylene,

R6 is COOR61, where

R61 is 1-4C-alkyl,

or the salts of this compound.

5. A compound of the formula I as claimed in claim 1 for use in the treatment of diseases.

6. A medicament comprising at least one compound of the formula I as claimed in claim 1 together with pharmaceutical auxiliaries and/or excipients.

7. The use of compounds of the formula I as claimed in claim 1 for the production of medicaments for the treatment of airway disorders.

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D221/12 A61K31/435 C07D491/04 //(C07D491/04,317:00,
221:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BIJAN PRASUN DAS ET AL: "Quinolizine derivatives: Part I Role of different cyclising agents in the formation of some ketoquinolizine derivatives" INDIAN JOURNAL OF CHEMISTRY, vol. 7, no. 2, 1969, pages 674-677, XP002048786 cited in the application see page 675, compound II, page 676 ---	1-3
A	DE 11 99 773 B (F. HOFFMANN-LA ROCHE & CO A.G.) 2 September 1965 cited in the application see the whole document --- -/--	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier document but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

11 November 1998

Date of mailing of the international search report

18/11/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Henry, J

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/04476

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 045 171 A (PFIZER INC.) 3 February 1982 see page 5 - page 10, line 5 see page 32, line 28 - page 39	1,5-7

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 1199773	B	NONE	
EP 0045171	A	03-02-1982	
		US 4309545 A	05-01-1982
		AR 230988 A	31-08-1984
		AT 11775 T	15-02-1985
		AU 525944 B	09-12-1982
		AU 7343481 A	06-05-1982
		CA 1178274 A	20-11-1984
		CS 228543 B	14-05-1984
		CS 228515 B	14-05-1984
		DD 202012 A	24-08-1983
		DK 334581 A	29-01-1982
		EG 15535 A	30-09-1986
		FI 812338 A	29-01-1982
		GR 75283 A	13-07-1984
		IE 51426 B	24-12-1986
		IN 158867 A	07-02-1987
		JP 57112372 A	13-07-1982
		PT 73440 B	20-06-1983
		SU 1217254 A	07-03-1986
		US 4351833 A	28-09-1982
		ZA 8105128 A	28-07-1982